



Correction to: Efficacy and Safety of Sublingual Fentanyl Tablets in Breakthrough Cancer Pain Management According to Cancer Stage and Background Opioid Medication

Jordi Guitart¹ · María Isabel Vargas² · Vicente De Sanctis³ · Jordi Folch¹ · Rafael Salazar⁴ · José Fuentes⁵ · Joan Coma⁶ · Julia Ferreras⁷ · Jordi Moya⁸ · Albert Tomás⁹ · Pere Estivill² · Francisco Rodelas⁴ · Antonio Javier Jiménez¹⁰ · Almudena Sanz¹⁰

Published online: 9 July 2018
© The Author(s) 2018

Correction to: Drugs R D (2018) 18:119–128
<https://doi.org/10.1007/s40268-018-0231-2>

In the Original Publication, Table 1 has error in the values. The values in the second column (LAC) and third column (Metastatic cancer) need to be swapped around.

Table 1 which currently reads:

Table 1 Patient demographics for the analyzed subgroups	Characteristic	LAC	Metastatic cancer	Fentanyl	Oxycodone/naloxone
	Age (years)	22 (70.4 ± 13.2)	54 (70.2 ± 11.9)	27 (67.7 ± 12.3)	24 (72.5 ± 10.5)
	Sex				
	Male	8 (34.8)	21 (36.2)	15 (51.7)	15 (62.5)
	Female	15 (65.2)	37 (63.8)	14 (48.3)	9 (37.5)
	Weight (kg)	20 (65.1 ± 13.7)	53 (72.1 ± 13.9)	26 (68.0 ± 12.4)	21 (73.7 ± 14.7)

Data are presented as *n* (%) or *N* (mean ± standard deviation)
LAC locally advanced cancer

The original article can be found online at <https://doi.org/10.1007/s40268-018-0231-2>.

✉ Jordi Guitart
jordi.guitart@hospitalplato.com

¹ Department of Anesthesiology, Hospital Plató, C/Plató 21, 08006 Barcelona, Spain

² Department of Anesthesiology, Parc Sanitari Sant Joan de Deu, Barcelona, Spain

³ Pain Unit, Department of Anesthesiology, Hospital Universitari Sagrat Cor, Barcelona, Spain

⁴ Department of Anesthesiology, Hospital Comarcal d’Inca, Palma de Mallorca, Spain

⁵ Department of Anesthesiology, Pius Hospital de Valls, Tarragona, Spain

⁶ Department of Anesthesiology, Hospital General de l’Hospitalet, Barcelona, Spain

⁷ Pain Unit, Department of Anesthesiology, Hospital Residència Sant Camil, Barcelona, Spain

⁸ Pain Unit, Department of Anesthesiology, Hospital Mateu Orfila, Menorca, Spain

⁹ Pain Unit, Department of Anesthesiology, Fundació Hospital Sant Bernabé, Barcelona, Spain

¹⁰ Kyowa Kirin Farmacéutica SLU, Madrid, Spain

Should read:

Table 1 Patient demographics for the analyzed subgroups

Characteristic	LAC	Metastatic cancer	Fentanyl	Oxycodone/naloxone
Age (years)	54 (70.2 ± 11.9)	22 (70.4 ± 13.2)	27 (67.7 ± 12.3)	24 (72.5 ± 10.5)
Sex				
Male	21 (36.2)	8 (34.8)	15 (51.7)	15 (62.5)
Female	37 (63.8)	15 (65.2)	14 (48.3)	9 (37.5)
Weight (kg)	53 (72.1 ± 13.9)	20 (65.1 ± 13.7)	26 (68.0 ± 12.4)	21 (73.7 ± 14.7)

Data are presented as *n* (%) or *N* (mean ± standard deviation)

LAC locally advanced cancer

Open Access This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.