

Association between subclinical atherosclerosis burden and unrecognized myocardial infarction detected by cardiac magnetic resonance in middle-aged low-risk adults

Leticia Fernández-Friera^{1,2,3,4,*†}, Ana García-Alvarez^{1,3,5†}, Belen Oliva¹,
Inés García-Lunar^{1,3,6}, Iris García¹, Andrea Moreno-Arciniegas^{1,7},
Sandra Gómez-Talavera^{1,7}, Cristina Pérez-Herreras⁸, Javier Sánchez-González⁹,
Vicente Martínez de Vega¹⁰, Xavier Rossello^{1,11}, Héctor Bueno^{1,3,12},
Antonio Fernández-Ortiz^{1,3,13}, Borja Ibañez^{1,3,7}, Javier Sanz^{1,14},
and Valentín Fuster^{1,14*}

¹Centro Nacional de Investigaciones Cardiovasculares (CNIC), Melchor Fernández Almagro 3, 28029, Madrid, Spain; ²HM CIEC MADRID (Centro Integral de Enfermedades Cardiovasculares), Hospital Universitario HM Montepíncipe, HM Hospitales, Av. de Montepíncipe, 25, 28660 Boadilla del Monte, Madrid, Spain; ³CIBERCV, Madrid, Spain; ⁴Universidad Camilo Jose Cela, Castillo de Alarcón, 49, 28692 Villafranca del Castillo, Madrid, Spain; ⁵Cardiology Department, Hospital Clinic Barcelona-IDIBAPS, Universitat de Barcelona, Barcelona, Spain; ⁶Cardiology Department, Hospital Universitario La Moraleja, Madrid, Spain; ⁷Cardiology Department, IIS-Hospital Universitario Fundación Jiménez Díaz, Av. de los Reyes Católicos 2, 28040 Madrid, Spain; ⁸Banco de Santander, Madrid, Spain; ⁹Philips Healthcare Iberia, Spain; ¹⁰Cardiology Department, Hospital Universitario Quirón, Madrid, Spain; ¹¹Cardiology Department, Hospital Universitari Son Espases-IDISBA, Palma de Mallorca, Spain; ¹²Cardiology Department, Hospital Universitario 12 de Octubre, and i+12 Research Institute, Madrid, Spain; ¹³Cardiology Department, Hospital Clínico San Carlos, Universidad Complutense, IdISSC, Madrid, Spain; and ¹⁴Mount Sinai Fuster Heart Hospital, New York, NY, USA

Received 22 July 2023; revised 13 November 2023; accepted 6 February 2024; online publish-ahead-of-print 1 March 2024

Aims

Evidence on the association between subclinical atherosclerosis (SA) and cardiovascular (CV) events in low-risk populations is scant. To study the association between SA burden and an ischaemic scar (IS), identified by cardiac magnetic resonance (CMR), as a surrogate of CV endpoint, in a low-risk population.

Methods and results

A cohort of 712 asymptomatic middle-aged individuals from the Progression of Early SA (PESA-CNIC-Santander) study (median age 51 years, 84% male, median SCORE2 3.37) were evaluated on enrolment and at 3-year follow-up with 2D/3D vascular ultrasound (VUS) and coronary artery calcification scoring (CACS). A cardiac magnetic study (CMR) was subsequently performed and IS defined as the presence of subendocardial or transmural late gadolinium enhancement (LGE). On CMR, 132 (19.1%) participants had positive LGE, and IS was identified in 20 (2.9%) participants. Individuals with IS had significantly higher SCORE2 at baseline and higher CACS and peripheral SA burden (number of plaques by 2DVUS and plaque volume by 3DVUS) at both SA evaluations. High CACS and peripheral SA (number of plaques) burden were independently associated with the presence of IS, after adjusting for SCORE2 [OR for 3rd tertile, 8.31; 95% confidence interval (CI) 2.85–24.2; $P < 0.001$; and 2.77; 95% CI, 1.02–7.51; $P = 0.045$, respectively] and provided significant incremental diagnostic value over SCORE2.

Conclusion

In a low-risk middle-aged population, SA burden (CAC and peripheral plaques) was independently associated with a higher prevalence of IS identified by CMR. These findings reinforce the value of SA evaluation to early implement preventive measures.

Clinical Trial Registration

Progression of Early Subclinical Atherosclerosis (PESA) Study Identifier: NCT01410318.

* Corresponding author. E-mail: vfuster@cnic.es (V.F.); E-mail: lfernandezfrierahmospitales.com (L.F.F.)

† These authors contributed equally to this work.

© The Author(s) 2024. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.




Figure 1 Prevalence of IS according to CAC, number of plaques by 2DVUS, and plaque volume by 3DVUS. CAC, Coronary artery calcification; MI, Myocardial infarction.

Table 3 CMR findings according to the presence of CAC or higher peripheral atherosclerosis burden by 2DVUS

	CAC		P	Peripheral atherosclerosis		P
	Absence (n = 239)	Presence (n = 468)		<Highest tertile (<8 plaques) (n = 505)	≥Highest tertile (≥8 plaques) (n = 183)	
LVEDVI (mL/m ²)	80.0 [71.8–88.9]	83.1 [73.9–91.2]	0.011	81.8 [73.2–90.8]	81.9 [73.3–91.1]	0.568
LVESVI (mL/m ²)	31.1 [26.3–36.1]	32.8 [27.8–37.9]	0.002	32.1 [27.1–37.4]	32.4 [27.4–37.5]	0.653
LVEF (%)	61.1 [58.0–64.6]	60.0 [56.8–63.6]	0.015	60.8 [57.1–63.8]	60.2 [56.9–64.2]	0.672
LVMI (g/m ²)	45.8 [39.4–53.2]	49.7 [44.1–55.9]	<0.001	48.4 [42.2–54.8]	49.0 [43.5–56.1]	0.130
Wall motion abnormalities	8 (3.3)	31 (6.6)	0.071	25 (4.9)	15 (8.2)	0.108
RVEDVI (mL/m ²)	81.2 [72.3–92.4]	83.8 [75.0–94.5]	0.024	83.7 [74.7–94.3]	81.4 [73.1–93.3]	0.184
RVESVI (mL/m ²)	31.8 [26.5–38.6]	34.4 [28.8–39.9]	0.003	34.4 [28.5–39.9]	32.7 [27.3–39.2]	0.114
RVEF (%)	60.2 [56.3–63.7]	59.0 [56.0–62.6]	0.011	59.3 [56–63]	59.7 [57–63.2]	0.168
LGE	32 (13.8)	98 (21.5)	0.016	82 (16.7)	44 (24.9)	0.017
Ischaemic	2 (0.9)	17 (3.7)	0.031	8 (1.6)	12 (6.8)	0.001
Non-ischaemic	30 (13.0)	81 (17.8)	0.108	74 (15.1)	32 (18.1)	0.348

CAC, coronary calcium score; LVEDVI, indexed left ventricle end-diastolic volume; LVESVI, indexed left ventricle end-systolic volume; LVEF, left ventricle ejection fraction; LVMI, indexed left ventricle mass; RVEDVI, indexed right ventricle end-diastolic volume; RVESVI, indexed right ventricle end-systolic volume; RVEF, left ventricle ejection fraction; LGE, late gadolinium enhancement. Significant P value (less than 0.05) in bold numbers.

a radiation-free technique, and suitable to be repeated in follow-up evaluations. Intriguingly, although both peripheral SA measures provided significant value, the number of plaques but not the 3DVUS plaque volume remained significantly associated with IS after adjusting by CV risk factors. The most plausible explanation for this finding is that plaque volume has been demonstrated to be very closely related to CV risk factors¹⁷ and because a higher number of plaques along the carotid and femoral arteries reflects a diffuse stage of atherosclerosis. The comparable results we obtained for CAC and VUS contrast with previous studies demonstrating the superiority of CAC over other measurements of peripheral SA, like carotid intima-media thickness, probably because VUS has demonstrated to better predict CV events than intima-media thickness,^{6,34} which is no longer recommended for CV risk assessment because it rather reflects hypertensive or age-related changes.^{1,35} In addition, in our cohort VUS examination was multi-territorial since it included both carotid, femoral arteries, and abdominal aorta. This approach has demonstrated diagnostic value,

particularly VUS of femoral arteries, in low-risk populations and the ability to increase the prediction of cardiovascular events compared with the measurement from a single site.⁶ In the BiImage study,³ which included ~6000 asymptomatic adults (mean age of 68.9 years), the impact of peripheral SA (as assessed by VUS of both carotid arteries) and CAC was cumulative suggesting that both techniques may be complementary in estimating CV risk. We obtained similar results using IS on CMR as an outcome instead of clinical events, due to the young and low-risk PESA population. In this line, scarce information exists regarding associations between peripheral atherosclerotic plaques and IS determined by CMR. In the study by Barbier et al, the presence of ≥50% arterial stenosis at whole-body magnetic resonance angiography was not associated with the presence of unrecognized MI.³⁶ This finding might be related to the different methods to assess atherosclerosis (severity of stenosis vs. plaque burden) and the larger diagnostic value of a measurement of the extension of disease than the presence of 'focal' narrowing.

All other authors have reported that they have no relationships to disclose relevant to the content of this paper.

Funding

The PESA study is co-funded equally by the Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain, and Banco Santander, Madrid, Spain. The study also receives funding from the Instituto de Salud Carlos III (ISCIII, PI15/02019) and the European Regional Development Fund (ERDF) 'Una manera de hacer Europa'. The CNIC is supported by the ISCIII, the Ministerio de Ciencia e Innovación and the Pro CNIC Foundation, and is a Severo Ochoa Center of Excellence (CEX2020-001041-S).

Data availability

It is feasible as far as it is required.

References

- Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Back M et al. 2021 ESC guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2021; **42**:3227–337.
- Peng AWW, Dardari ZA, Blumenthal RS, Dzay O, Obisesan OH, Iftekhar Uddin SM et al. Very high coronary artery calcium (≥ 1000) and association with cardiovascular disease events, non-cardiovascular disease outcomes, and mortality: results from MESA. *Circulation* 2021; **143**:1571–83.
- Baber U, Mehran R, Sartori S, Schoos MM, Sillesen H, Muntendam P et al. Prevalence, impact, and predictive value of detecting subclinical coronary and carotid atherosclerosis in asymptomatic adults: the BiImage study. *J Am Coll Cardiol* 2015; **65**:1065–74.
- Blaaha MJ, Whelton SP, Al Rifai M, Dardari Z, Shaw LJ, Al-Mallah MH et al. Comparing risk scores in the prediction of coronary and cardiovascular deaths: coronary artery calcium consortium. *JACC Cardiovasc Imaging* 2021; **14**:411–21.
- Wright JD, Folsom AR, Coresh J, Sharrett AR, Couper D, Wagenknecht LE et al. The ARIC (atherosclerosis risk in communities) study: JACC focus seminar 3/8. *J Am Coll Cardiol* 2021; **77**:2939–59.
- Nicolaidis AN, Panayiotou AG, Griffin M, Tyllis T, Bond D, Georgiou N et al. Arterial ultrasound testing to predict atherosclerotic cardiovascular events. *J Am Coll Cardiol* 2022; **79**:1969–82.
- Redfield MM. Heart failure—an epidemic of uncertain proportions. *N Engl J Med* 2002; **347**:1442–4.
- Turkbeey EB, Nacif MS, Guo M, McClelland RL, Teixeira PB, Bild DE et al. Prevalence and correlates of myocardial scar in a US cohort. *JAMA* 2015; **314**:1945–54.
- de Torbal A, Boersma E, Kors JA, van Herpen G, Deckers JW, van der Kuip DA et al. Incidence of recognized and unrecognized myocardial infarction in men and women aged 55 and older: the Rotterdam study. *Eur Heart J* 2006; **27**:729–36.
- Yang Y, Li W, Zhu H, Pan XF, Hu Y, Arnott C et al. Prognosis of unrecognized myocardial infarction determined by electrocardiography or cardiac magnetic resonance imaging: systematic review and meta-analysis. *BMJ* 2020; **369**:m1184.
- Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA et al. Fourth universal definition of myocardial infarction (2018). *Eur Heart J* 2019; **40**:237–69.
- Fernandez-Ortiz A, Jimenez-Borreguero LJ, Penalvo JL, Ordovas JM, Mocoora A, Fernandez-Friera L et al. The progression and early detection of subclinical atherosclerosis (PESA) study: rationale and design. *Am Heart J* 2013; **166**:990–8.
- Ibanez B, Fernandez-Ortiz A, Fernandez-Friera L, Garcia-Lunar I, Andres V, Fuster V. Progression of early subclinical atherosclerosis (PESA) study: JACC focus seminar 7/8. *J Am Coll Cardiol* 2021; **78**:156–79.
- collaboration S-OwgaECr. SCORE2-OP risk prediction algorithms: estimating incident cardiovascular event risk in older persons in four geographical risk regions. *Eur Heart J* 2021; **42**:2455–67.
- Cerqueira MD, Weissman NJ, Dilsizian V, Jacobs AK, Kaul S, Laskey WK et al. Standardized myocardial segmentation and nomenclature for tomographic imaging of the heart. A statement for healthcare professionals from the cardiac imaging committee of the council on clinical cardiology of the American heart association. *Circulation* 2002; **105**:539–42.
- Shanbhag SM, Greve AM, Aspelund T, Schelbert EB, Cao JJ, Danielsen R et al. Prevalence and prognosis of ischaemic and non-ischaemic myocardial fibrosis in older adults. *Eur Heart J* 2019; **40**:529–38.
- Lopez-Melgar B, Fernandez-Friera L, Oliva B, Garcia-Ruiz JM, Penalvo JL, Gomez-Talavera S et al. Subclinical atherosclerosis burden by 3D ultrasound in mid-life: the PESA study. *J Am Coll Cardiol* 2017; **70**:301–13.
- Fernandez-Friera L, Penalvo JL, Fernandez-Ortiz A, Ibanez B, Lopez-Melgar B, Laclaustra M et al. Prevalence, vascular distribution, and multiterritorial extent of subclinical atherosclerosis in a middle-aged cohort: the PESA (progression of early subclinical atherosclerosis) study. *Circulation* 2015; **131**:2104–13.
- Agatston AS, Janowitz WR, Hildner FJ, Zusmer NR, Viamonte M Jr, Detrano R. Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol* 1990; **15**:827–32.
- Lopez-Melgar B, Fernandez-Friera L, Oliva B, Garcia-Ruiz JM, Sanchez-Cabo F, Bueno H et al. Short-Term progression of multiterritorial subclinical atherosclerosis. *J Am Coll Cardiol* 2020; **75**:1617–27.
- Kwong RY, Chan AK, Brown KA, Chan CW, Reynolds HG, Tsang S et al. Impact of unrecognized myocardial scar detected by cardiac magnetic resonance imaging on event-free survival in patients presenting with signs or symptoms of coronary artery disease. *Circulation* 2006; **113**:2733–43.
- Barbier CE, Bjerner T, Johansson L, Lind L, Ahlstrom H. Myocardial scars more frequent than expected: magnetic resonance imaging detects potential risk group. *J Am Coll Cardiol* 2006; **48**:765–71.
- Weir-McCall JR, Fitzgerald K, Papagiorcopulo CJ, Gandy SJ, Lambert M, Belch JFF et al. Prevalence of unrecognized myocardial infarction in a low-intermediate risk asymptomatic cohort and its relation to systemic atherosclerosis. *Eur Heart J Cardiovasc Imaging* 2017; **18**:657–62.
- Yang Z, Fu H, Li H, Wang JR, Xu HY, Xie LJ et al. Late gadolinium enhancement is a risk factor for major adverse cardiac events in unrecognized myocardial infarction without apparent symptoms: a meta-analysis. *Clin Radiol* 2021; **76**:79 e1–79.e11.
- Kim HW, Klem I, Shah DJ, Wu E, Meyers SN, Parker MA et al. Unrecognized non-Q-wave myocardial infarction: prevalence and prognostic significance in patients with suspected coronary disease. *PLoS Med* 2009; **6**:e1000057.
- Goehde SC, Hunold P, Vogt FM, Ajaaj W, Goyen M, Herborn CU et al. Full-body cardiovascular and tumor MRI for early detection of disease: feasibility and initial experience in 298 subjects. *AJR Am J Roentgenol* 2005; **184**:598–611.
- Blaaha MJ, DeFilippis AP. Multi-Ethnic study of atherosclerosis (MESA): JACC focus seminar 5/8. *J Am Coll Cardiol* 2021; **77**:3195–216.
- Hofman A, Breteler MM, van Duijn CM, Krestin GP, Pols HA, Stricker BH et al. The rotterdam study: objectives and design update. *Eur J Epidemiol* 2007; **22**:819–29.
- Erbel R, Eisele L, Moebus S, Dragano N, Möhlenkamp S, Bauer M et al. [The heinz nixdorf recall study]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2012; **55**(6–7):809–15.
- Edvardsson T, Detrano R, Rosen BD, Carr JJ, Liu K, Lai S et al. Coronary artery atherosclerosis is related to reduced regional left ventricular function in individuals without history of clinical cardiovascular disease: the multiethnic study of atherosclerosis. *Arterioscler Thromb Vasc Biol* 2006; **26**:206–11.
- Bakhshi H, Ambale-Venkatesh B, Yang X, Ostovaneh MR, Wu CO, Budoff M et al. Progression of coronary artery calcium and incident heart failure: the multi-ethnic study of atherosclerosis. *J Am Heart Assoc* 2017; **6**:e005253.
- Yared GS, Moreira HT, Ambale-Venkatesh B, Vasconcelos HD, Nwabuo CC, Ostovaneh MR et al. Coronary artery calcium from early adulthood to middle age and left ventricular structure and function. *Circ Cardiovasc Imaging* 2019; **12**:e009228.
- Schelbert EB, Cao JJ, Sigurdsson S, Aspelund T, Kellman P, Aletras AH et al. Prevalence and prognosis of unrecognized myocardial infarction determined by cardiac magnetic resonance in older adults. *JAMA* 2012; **308**:890–6.
- Poredos P, Cifkova R, Marie Maier JA, Nemcsik J, Sabovic M, Jug B et al. Preclinical atherosclerosis and cardiovascular events: do we have a consensus about the role of pre-clinical atherosclerosis in the prediction of cardiovascular events? *Atherosclerosis* 2022; **348**:25–35.
- Mach F, Baigent C, Catapano AL, Koskinas KC, Casula M, Badimon L et al. 2019 ESC/EAS guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J* 2020; **41**:111–88.
- Ebeling Barbier C, Bjerner T, Hansen T, Andersson J, Lind L, Hulthe J et al. Clinically unrecognized myocardial infarction detected at MR imaging may not be associated with atherosclerosis. *Radiology* 2007; **245**:103–10.